

09870483, 060101

09870483, 060101

09870483, 060101

09870483, 060101

09870483, 060101

09870483, 060101

In these methods, hydrolysis is performed after the isolation of optically active cyanohydrin, or in consideration of efficiency, hydrolysis is performed without the removal of the reaction solvent used in the production process of the optically active cyanohydrin.

The known methods for synthesizing optically active α -hydroxycarboxylic acid by hydrolysis of optically active cyanohydrin include, for example, a method described in *Synthesis*, July 1990, 575-578 and a method described in *Tetrahedron Letters*, 32, 2605-2608 (1991), wherein reaction with concentrated hydrochloric acid at room temperature is followed by reaction at reflux temperature.

With regard to these methods, however, since a large amount of hydrochloric acid, about 30 times the amount of cyanohydrin by molar ratio is used, industrial application of these methods involves many disadvantages such as the high-cost raw materials, low productivity related to the necessity of a huge reactor, a large amount of waste liquid and so on. Furthermore, for the reason that generally α -hydroxycarboxylic acid of interest is highly soluble in water, in a case where, after reaction with a large amount of acid, the substance is purified by operations such as extraction or crystallization, a large amount of substance may disadvantageously remain in aqueous solution, having low yield.

Still more, where inappropriate reaction conditions are applied to a reaction, a side reaction or racemization reaction may often occur, resulting in a low yield and low optical purity of α -hydroxycarboxylic acid.

OBJECTS AND SUMMARY OF THE INVENTION

09370483-060101 TOTAL 2949/2950

The object of the first aspect of the present invention is to provide a method for producing high purity α -hydroxycarboxylic acid.

The object of the second and third aspects of the present invention is to provide an industrially advantageous method for producing optically active α -hydroxycarboxylic acid.

As a result of thorough studies by the present inventors directed toward the object of the first aspect above, we have found that the object can be achieved by hydrolyzing cyanohydrin in the presence of a hydrocarbon solvent, thereby completing the first aspect of the present invention.

That is to say, the first aspect of the present invention comprises the following features of invention.

- (1) A method for producing α -hydroxycarboxylic acid, which comprises hydrolyzing cyanohydrin in the presence of a hydrocarbon solvent.
- (2) The method for producing α -hydroxycarboxylic acid according to (1) above, which comprises separating and removing the hydrocarbon solvent phase from a reaction solution after hydrolysis reaction.
- (3) The method for producing α -hydroxycarboxylic acid according to (1) or (2) above, wherein the hydrolysis reaction is carried out using mineral acid.

Moreover, as a result of thorough studies by the present inventors to improve the yield and optical purity of optically active α -hydroxycarboxylic acid, we have found that, in a case where an alcoholic solvent, an ester solvent, an ethereal solvent, a carboxylic solvent or a hydrocarbon-based solvent is used as a reaction solvent in the

production process of optically active cyanohydrin, followed by the removal of the solvent and hydrolysis, both the yield and optical purity of optically active α -hydroxycarboxylic acid are improved, compared with a case where hydrolysis is performed without removing the solvent; thereby completing the second aspect of the present invention.

That is to say, the second aspect of the present invention comprises the following features of invention.

- (1) A method for producing optically active α -hydroxycarboxylic acid, which comprises: producing optically active cyanohydrin by performing a reaction between a carbonyl compound and hydrogen cyanide, using a solvent comprising at least one organic solvent selected from a group consisting of an alcoholic solvent, an ester solvent, an ethereal solvent and a carboxylic solvent; removing the above organic solvent from the above reaction solvent; and hydrolyzing the remaining reaction mixture without isolating optically active cyanohydrin.
- (2) The method for producing optically active α -hydroxycarboxylic acid according to (1) above, wherein the amount of the above organic solvent in the reaction mixture subjected to hydrolysis is less than 10 weight %.
- (3) The method for producing optically active α -hydroxycarboxylic acid according to (1) or (2) above, wherein the hydrolysis reaction is carried out using mineral acid.

As a result of further thorough studies by the present inventors directed toward the object of the third aspect of the present invention, we have found that the object can be achieved by hydrolyzing optically active cyanohydrin, using at most 10 equivalents of mineral acid relative to the above optically active cyanohydrin under the condition that maximum temperature on reaction time is 90°C or less; thereby

completing the third aspect of the present invention.

That is to say, the third aspect of the present invention comprises the following feature of invention.

(1) A method for producing optically active α -hydroxycarboxylic acid, which comprises hydrolyzing optically active cyanohydrin, using at most 10 equivalents of mineral acid relative to said optically active cyanohydrin under the condition that maximum temperature on reaction time is 90°C or less.

Furthermore, the present inventors have found that the crystal with high packing density of optically active α -hydroxycarboxylic acid can be obtained by crystallizing optically active α -hydroxycarboxylic acid in an aqueous solution, thereby completing a fourth aspect of the present invention.

That is to say, the fourth aspect of the present invention comprises the following features of invention.

(1) A method for producing optically active crystalline α -hydroxycarboxylic acid, which comprises crystallizing optically active α -hydroxycarboxylic acid in an aqueous solution.

(2) The method for producing optically active crystalline α -hydroxycarboxylic acid according to (1) above, which comprises crystallizing optically active α -hydroxycarboxylic acid in the presence of a non-miscible organic solvent.

(3) An optically active crystalline chloromandelic acid, which is obtained by the production method according to (1) or (2) above.


09870483-060101

(4) A method for producing optically active crystalline α -hydroxycarboxylic acid, which comprises crystallizing the optically active α -hydroxycarboxylic acid obtained by the method according to the first, second or third aspect of the present invention, in an aqueous solution.

The method of the first aspect of the present invention is to produce α -hydroxycarboxylic acid by hydrolyzing cyanohydrin in the presence of a hydrocarbon solvent and converting the cyano group of the above cyanohydrin into a carboxyl group, and according to the present method, it becomes possible to easily remove colored substances and by-products generated during the hydrolysis reaction, resulting in the simple production of high purity α -hydroxycarboxylic acid.

In the first aspect of the present invention, cyanohydrin used as a material for the present invention is not particularly limited, as long as it has, in a molecule thereof, at least one pair consisting of a hydroxyl group and a cyano group which bind to an identical carbon atom.

An example of cyanohydrin used for the present invention includes the compound shown in the following formula (I):



wherein

R^1 and R^2 may be different from or identical to each other, independently representing a

hydrogen atom, a halogen atom, an amino group, an amino group mono- or di-substituted with a monovalent hydrocarbon group containing at most 14 carbon atoms, a mercapto group or a monovalent hydrocarbon group containing at most 22 carbon atoms, in the above hydrocarbon group, each of $-\text{CH}_2-$ and CH_2 in $-\text{CH}_3$ may be substituted with a carbonyl group, a sulfonyl group, $-\text{O}-$ or $-\text{S}-$, $=\text{CH}_2$ may be substituted with $=\text{O}$ or $=\text{S}$; or C-H in $-\text{CH}_2$, C-H in $-\text{CH}_3$, C-H in $>\text{CH}-$, C-H in $=\text{CH}-$ and C-H in $=\text{CH}_2$ may be substituted with N or C-halogen, or R^1 and R^2 may together form a divalent group.

The monovalent hydrocarbon group containing at most 22 carbon atoms in the above formula (I) includes a linear or branched chain hydrocarbon group, a monocyclic hydrocarbon group with or without side chain, a polycyclic hydrocarbon group with or without side chain, a spiro hydrocarbon group with or without side chain, a ring-assembled structural hydrocarbon group with or without side chain, or a chain hydrocarbon group with the above cyclic-hydrocarbon. It includes any saturated or unsaturated hydrocarbon group, with the exception that unsaturated hydrocarbon groups having an allene structure ($\text{C}=\text{C}=\text{C}$) are excluded. The linear or branched chain hydrocarbon group includes, for example, saturated chain hydrocarbon groups such as a linear alkyl group containing at least 1 carbon atom and a branched alkyl group containing at least 3 carbon atoms; unsaturated chain hydrocarbon groups such as a linear alkenyl group containing at least 2 carbon atoms, a branched alkenyl group containing at least 3 carbon atoms, a linear alkynyl group containing at least 3 carbon atoms, a branched alkynyl group containing at least 4 carbon atoms, a linear alkadienyl group containing at least 4 carbon atoms and a branched alkadienyl group containing at least 5 carbon atoms. The monocyclic hydrocarbon group includes, for example, saturated monocyclic hydrocarbon groups such as a cycloalkyl group without side chain which contains at least 3 carbon atoms and a cycloalkyl group with side chain which contains at least 4 carbon atoms in total; unsaturated monocyclic hydrocarbon groups such as a cycloalkenyl group without side chain which contains at least 4 carbon atoms,

a cycloalkynyl group with side chain which contains at least 5 carbon atoms in total, a cycloalkadienyl group without side chain which contains at least 5 carbon atoms and a cycloalkadienyl group with side chain which contains at least 6 carbon atoms in total. The unsaturated monocyclic or polycyclic hydrocarbon group includes an aromatic hydrocarbon group including: an aromatic group without side chain which contains 6 to 22 carbon atoms in total such as a phenyl group, a 1-naphthyl group, a 2-naphthyl group and a 9-anthryl group; an aromatic group with side chain which contains at least 7 carbon atoms in total; a phenylphenyl group containing 12 carbon atoms and a phenylphenyl group with side chain which contains at least 13 carbon atoms in total, which are also included in a ring-assembled structural hydrocarbon group. The polycyclic hydrocarbon group includes a condensed cyclic hydrocarbon group without side chain which contains at least 6 carbon atoms, a condensed cyclic hydrocarbon group with side chain which contains at least 7 carbon atoms in total, a bridged cyclic hydrocarbon group without side chain which contains at least 7 carbon atoms, a bridged cyclic hydrocarbon group with side chain which contains at least 8 carbon atoms in total, a spiro hydrocarbon group without side chain which contains at least 9 carbon atoms in total and a spiro hydrocarbon group with side chain which contains at least 10 carbon atoms in total. In addition, the above condensed cyclic hydrocarbon group without side chain includes those which contain at least 9 carbon atoms in total when one of its condensed rings is a benzene ring, and the above condensed cyclic hydrocarbon group with side chain includes those which contain at least 10 carbon atoms in total when one of its condensed rings is benzene ring. The ring-assembled structural hydrocarbon group includes a cycloalkyl-cycloalkyl group without side chain which contains at least 6 carbon atoms in total, a cycloalkyl-cycloalkyl group with side chain which contains at least 7 carbon atoms in total, a cycloalkylidene-cycloalkyl group without side chain which contains at least 6 carbon atoms in total and a cycloalkylidene-cycloalkyl group with side chain which contains at least 7 carbon atoms in total. "A cyclic hydrocarbon with side chain" in these cyclic hydrocarbons, corresponds to one having a chain hydrocarbon group attached to its ring. Such chain hydrocarbon group having a cyclic

hydrocarbon includes a linear alkyl group which is substituted with an aromatic group without side chain and contains at least 7 carbon atoms in total, a linear alkyl group which is substituted with an aromatic group with side chain and contains at least 8 carbon atoms in total, a branched alkyl group which is substituted with an aromatic group without side chain and contains at least 9 carbon atoms in total, a branched alkyl group which is substituted with an aromatic group with side chain and contains at least 10 carbon atoms in total, a linear alkenyl group which is substituted with an aromatic group without side chain and contains at least 8 carbon atoms in total, a linear alkenyl group which is substituted with an aromatic group with side chain and contains at least 9 carbon atoms in total, a branched alkenyl group which is substituted with an aromatic group without side chain and contains at least 9 carbon atoms in total, a branched alkenyl group which is substituted with an aromatic group with side chain and contains at least 10 carbon atoms in total, a linear alkynyl group which is substituted with an aromatic group without side chain and contains at least 8 carbon atoms in total, a linear alkynyl group which is substituted with an aromatic group with side chain and contains at least 9 carbon atoms in total, a branched alkynyl group which is substituted with an aromatic group without side chain and contains at least 10 carbon atoms in total, a branched alkynyl group which is substituted with an aromatic group with side chain and contains at least 11 carbon atoms in total, a linear alkadienyl group which is substituted with an aromatic group without side chain and contains at least 10 carbon atoms in total, a linear alkadienyl group which is substituted with an aromatic group with side chain and contains at least 11 carbon atoms in total, a branched alkadienyl group which is substituted with an aromatic group without side chain and contains at least 11 carbon atoms in total, a branched alkadienyl group which is substituted with an aromatic group with side chain and contains at least 12 carbon atoms in total, a linear alkyl group which is substituted with a cycloalkyl group without side chain and contains at least 4 carbon atoms in total, a linear alkyl group which is substituted with a cycloalkyl group with side chain and contains at least 5 carbon atoms in total, a branched alkyl group which is substituted with a cycloalkyl group without side chain and contains at least 6 carbon

atoms in total, a branched alkyl group which is substituted with a cycloalkyl group with side chain and contains at least 7 carbon atoms in total, a linear alkenyl group which is substituted with a cycloalkyl group without side chain and contains at least 5 carbon atoms in total, a linear alkenyl group which is substituted with a cycloalkyl group with side chain and contains at least 6 carbon atoms in total, a branched alkenyl group which is substituted with a cycloalkyl group without side chain and contains at least 6 carbon atoms in total, a branched alkenyl group which is substituted with a cycloalkyl group with side chain and contains at least 7 carbon atoms in total, a linear alkynyl group which is substituted with a cycloalkyl group without side chain and contains at least 5 carbon atoms in total, a linear alkynyl group which is substituted with a cycloalkyl group with side chain and contains at least 6 carbon atoms in total, a branched alkynyl group which is substituted with a cycloalkyl group without side chain and contains at least 7 carbon atoms in total, a branched alkynyl group which is substituted with a cycloalkyl group with side chain and contains at least 8 carbon atoms in total, a branched alkadienyl group which is substituted with a cycloalkyl group without side chain and contains at least 8 carbon atoms in total, a branched alkadienyl group which is substituted with a cycloalkyl group with side chain and contains at least 9 carbon atoms in total.

Hereinafter, each of an aromatic group without side chain, an aromatic group with side chain, and a phenylphenyl group or a phenylphenyl group with side chain, is referred as an aryl group, and a linear or branched alkyl group substituted with the aryl group or groups is referred as an aralkyl group. Other cyclic hydrocarbon groups referring to both one having side chains on its ring and one having no side chains, are simply referred as, for example, cycloalkyl groups, unless otherwise specified. Also, chain hydrocarbon groups referring to both linear one and branched one are simply referred as, for example, alkyl groups.

When $-\text{CH}_2-$ in the above hydrocarbon group is substituted with a carbonyl

group, sulfonyl group, -O- or -S-, a ketone, sulfone, ether or thioether structure is introduced therein, respectively. When $-\text{CH}_2-$ in $-\text{CH}_3$ is substituted with a carbonyl group, -O- or -S-, it converts into a formyl (aldehyde) group, a hydroxyl group or a mercapto group, respectively. When a terminal $=\text{CH}_2$ is substituted with $=\text{O}$ or $=\text{S}$, a ketone or thioketone structure is introduced therein. When each C-H in $-\text{CH}_2-$ is substituted with N, it converts into -NH-. When C-H in $>\text{CH}-$ is substituted with N, it converts into $>\text{N}-$. When C-H in $=\text{CH}-$ is substituted with N, it converts into $=\text{N}-$. When C-H in a terminal $-\text{CH}_3$ is substituted with N, $-\text{NH}_2$ is introduced therein. When C-H in $=\text{CH}_2$ is substituted with N, it converts into $=\text{NH}$. Further, each C-H in $-\text{CH}_3$, $-\text{CH}_2-$, $=\text{CH}-$, $\equiv\text{CH}$ or $>\text{CH}-$ is substituted with a C-halogen, the carbon is substituted with a halogen atom. The substitution of carbon chains with -O-, -S- or N corresponds to oxa-, thia- or aza-substitution of the hydrocarbon group, respectively. For example, when these substitution take place in a ring carbon of the hydrocarbon ring, the hydrocarbon ring converts into a heterocyclic ring respectively containing oxygen, sulfur or nitrogen. The substitution of CH_2 and C-H in the hydrocarbon group may independently take place and it may further take place when CH_2 or C-H still remains on the carbon after the prior substitution. Further, the above substitutions may bring the conversion of $-\text{CH}_2-\text{CH}_3$ into $-\text{CO}-\text{O}-\text{H}$, a carboxylic acid structure.

A halogen atom, herein, refers to a fluorine atom, a chlorine atom, a bromine atom and an iodine atom, but a fluorine atom, a chlorine atom and a bromine atom are preferable.

Accordingly, the above hydrocarbon group may be selected from any chain hydrocarbon group and hydrocarbon group having ring-structured hydrocarbon group such as cyclic hydrocarbon groups, for example, saturated chain hydrocarbon groups such as linear or branched alkyl groups; unsaturated chain hydrocarbon groups such as linear or branched alkenyl groups, linear or branched alkynyl groups and linear or branched alkadienyl groups; saturated cyclic hydrocarbon groups such as cycloalkyl

groups; unsaturated cyclic hydrocarbon groups such as cycloalkenyl groups, cycloalkynyl groups and cycloalkadienyl groups; and aromatic hydrocarbon groups such as aryl groups, aralkyl groups and arylalkenyl groups.

In more detail, the linear or branched alkyl groups include a methyl group, an ethyl group, a propyl group, an isopropyl group, a butyl group, a 1-methylpropyl group, a pentyl group, a 1-methylbutyl group, a hexyl group, a 1-methylpentyl group, a heptyl group, a 1-methylhexyl group, a 1-ethylpentyl group, an octyl group, a nonyl group, a decyl group, an undecyl group, a dodecyl group, a tridecyl group, a tetradecyl group, a 2-methylpropyl group, a 2-methylbutyl group, a 3-methylbutyl group, a 2-methylpentyl group, a 3-methylpentyl group, a 4-methylpentyl group, a methylhexyl group, a methylheptyl group, a methyloctyl group, a methylnonyl group, a 1,1-dimethylethyl group, a 1,1-dimethylpropyl group, a 2,6-dimethylheptyl group, a 3,7-dimethyloctyl group and a 2-ethylhexyl group; cycloalkylalkyl groups include a cyclopentylmethyl group and a cyclohexylmethyl group; cycloalkyl groups include a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, a methylcyclopentyl group, a cyclohexyl group, a methylcyclohexyl group, a cycloheptyl group and a cyclooctyl group; and bicycloalkyl groups include a norbornyl group, a bicyclo [2.2.2] octyl group and an adamantyl group.

The linear or branched alkenyl groups include a vinyl group, an allyl group, a crotyl group (a 2-butenyl group) and an isopropenyl group (a 1-methylvinyl group); cycloalkenyl or cycloalkadienyl groups include a cyclopentenyl group, a cyclopentadienyl group, a cyclohexenyl group and a cyclohexadienyl group.

The linear or branched alkynyl groups include an ethynyl group, a propynyl group and a butynyl group. The aryl groups include a phenyl group, a 1-naphthyl group, a 2-naphthyl group, a 2-phenylphenyl group, a 3-phenylphenyl group, a 4-

phenylphenyl group, a 9-anthryl group, a methylphenyl group, a dimethylphenyl group, a trimethylphenyl group, an ethylphenyl group, a methylethylphenyl group, a diethylphenyl group, a propylphenyl group and a butylphenyl group.

The aralkyl groups include a benzyl group, a 1-naphthylmethyl group, a 2-naphthylmethyl group, a phenethyl group (a 2-phenylethyl group), a 1-phenylethyl group, a phenylpropyl group, a phenylbutyl group, a phenylpentyl group, a phenylhexyl group, a methylbenzyl group, a methylphenethyl group, a dimethylbenzyl group, a dimethylphenethyl group, a trimethylbenzyl group, an ethylbenzyl group and a diethylbenzyl group.

The arylalkenyl groups include a styryl group, a methylstyryl group, an ethylstyryl group, a dimethylstyryl group and a 3-phenyl-2-propenyl group.

The above hydrocarbon groups, in which the CH_2 group is substituted with a carbonyl group, a sulfonyl group, O or S, or the C-H group is substituted with N or C-halogen, include groups having one or more structures such as ketone, aldehyde, carboxylic acid, sulfone, ether, thioether, amine, alcohol, thiol, halogen and heterocycles (e.g. an oxygen-containing heterocycle, a sulfur-containing heterocycle, a nitrogen-containing heterocycle, etc.) The oxygen-containing heterocycle, sulfur-containing heterocycle and nitrogen-containing heterocycle correspond to cyclic hydrocarbon groups in which their ring carbon is substituted with oxygen, sulfur and nitrogen, respectively. These heterocycles may contain two or more heteroatoms.

These substituted hydrocarbon groups may include a ketone structure such as an acetylmethyl group and an acetylphenyl group; a sulfone structure such as a methanesulfonylmethyl group; an ether structure such as a methoxymethyl group, a

methoxyethyl group, an ethoxyethyl group, a methoxypropyl group, a butoxyethyl group, an ethoxyethoxyethyl group, a methoxyphenyl group, dimethoxyphenyl group and phenoxyethyl group; a thioether structure such as a methylthiomethyl group and a methylthiophenyl group; an amine structure such as an aminomethyl group, a 2-aminoethyl group, a 2-aminopropyl group, a 3-aminopropyl group, a 2,3-diaminopropyl group, a 2-aminobutyl group, a 3-aminobutyl group, a 4-aminobutyl group, a 2,3-diaminobutyl group, a 2,4-diaminobutyl group, a 3,4-diaminobutyl group, a 2,3,4-triaminobutyl group, a methylaminomethyl group, a dimethylaminomethyl group, a methylaminoethyl group, a propylaminomethyl group, a cyclopentylaminomethyl group, an aminophenyl group, a diaminophenyl group, an aminomethylphenyl group; an oxygen-containing heterocycle such as a tetrahydrofuran group, a tetrahydropyran group and a morpholethyl group; an oxygen-containing aromatic ring such as a furyl group, a furfuryl group, a benzofuryl group and a benzofurfuryl group; a sulfur-containing heterocycle such as a thienyl group; a nitrogen-containing aromatic ring such as a pyrrolyl group, an imidazolyl group, an oxazolyl group, a thiazolyl group, a pyridyl group, a pyrimidyl group, a pyridazinyl group, a pyrazinyl group, a tetrazinyl group, a quinolyl group, an isoquinolyl group, a pyridylmethyl group; an alcohol structure such as a 2-hydroxyethyl group, a 2-hydroxypropyl group, a 3-hydroxypropyl group, a 2,3-dihydroxypropyl group, a 2-hydroxybutyl group, a 3-hydroxybutyl group, a 4-hydroxybutyl group, a 2,3-dihydroxybutyl group, a 2,4-dihydroxybutyl group, a 3,4-dihydroxybutyl group, a 2,3,4-trihydroxybutyl group, a hydroxyphenyl group, a dihydroxyphenyl group, a hydroxymethylphenyl group and a hydroxyethylphenyl group; a thiol structure such as a 2-mercaptoethyl group, a 2-mercaptopropyl group, a 3-mercaptopropyl group, a 2,3-dimercaptopropyl group, a 2-mercaptobutyl group, a 3-mercaptobutyl group, a 4-mercaptobutyl group, a mercaptophenyl group; a halogenated hydrocarbon group such as a 2-chloroethyl group, a 2-chloropropyl group, a 3-chloropropyl group, a 2-chlorobutyl group, a 3-chlorobutyl group, a 4-chlorobutyl group,

a fluorophenyl group, a chlorophenyl group, a bromophenyl group, a difluorophenyl group, a dichlorophenyl group, a dibromophenyl group, a chlorofluorophenyl group, a trifluorophenyl group, a trichlorophenyl group, a fluoromethylphenyl group and a trifluoromethylphenyl group; a compound having both an amine structure and an alcohol structure such as a 2-amino-3-hydroxypropyl group, a 3-amino-2-hydroxypropyl group, a 2-amino-3-hydroxybutyl group, a 3-amino-2-hydroxybutyl group, a 2-amino-4-hydroxybutyl group, a 4-amino-2-hydroxybutyl group, a 3-amino-4-hydroxybutyl group, a 4-amino-3-hydroxybutyl group, a 2,4-diamino-3-hydroxybutyl group, a 3-amino-2,4-dihydroxybutyl group, a 2,3-diamino-4-hydroxybutyl group, a 4-amino-2,3-dihydroxybutyl group, a 3,4-diamino-2-hydroxybutyl group, a 2-amino-3,4-dihydroxybutyl group and an aminohydroxyphenyl group; a hydrocarbon group substituted with a halogen and a hydroxyl group such as a fluorohydroxyphenyl group, a chlorohydroxyphenyl group; and a carbon structure such as a carboxyphenyl group.

The cyanohydrin shown in the above formula (I) includes: 2-aryl-2-hydroxyacetonitrile such as mandelonitrile (2-hydroxy-2-phenylacetonitrile), 3-phenoxymandelonitrile (2-hydroxy-2-(3-phenoxyphenyl)acetonitrile), 4-methylmandelonitrile (2-hydroxy-2-(p-tolyl)acetonitrile), 2-chloromandelonitrile (2-(2-chlorophenyl)-2-hydroxyacetonitrile), 3-chloromandelonitrile (2-(3-chlorophenyl)-2-hydroxyacetonitrile), 4-chloromandelonitrile (2-(4-chlorophenyl)-2-hydroxyacetonitrile), 3-nitromandelonitrile (2-hydroxy-2-(3-nitrophenyl)acetonitrile), 3,4-methylenedioxymandelonitrile (2-hydroxy-2-(3,4-methylenedioxyphenyl)acetonitrile), 2,3-methylenedioxymandelonitrile (2-hydroxy-2-(2,3-methylenedioxyphenyl)acetonitrile), 2-benzyl-2-hydroxyacetonitrile and 2-(2-furyl)-2-hydroxyacetonitrile; 2-alkyl-2-hydroxyacetonitrile such as 2-hydroxy-2-methylacetonitrile, 2-hydroxy-2-propylacetonitrile, 2-hydroxy-2-isopropylacetonitrile, 2-butyl-2-hydroxyacetonitrile and 2-cyclohexyl-2-hydroxyacetonitrile; 2,2-dialkyl-2-hydroxyacetonitrile such as 2-ethyl-2-hydroxy-2-methylacetonitrile, 2-butyl-2-hydroxy-

2-methylacetonitrile, 2-hydroxy-2-methyl-2-propylacetonitrile, 2-hydroxy-2-isopropyl-2-methylacetonitrile, 2-hydroxy-2-methyl-2-pentylacetonitrile, 2-hydroxy-2-methyl-2-(2-methylpropyl)acetonitrile and 2-hydroxy-2-methyl-2-(3-methylbutyl)acetonitrile; 2-alkyl-2-alkenyl-2-hydroxyacetonitrile such as 2-hydroxy-2-methyl-2-(2-propenyl)acetonitrile and 2-(3-butenyl)-2-hydroxy-2-methylacetonitrile; 2-alkyl-2-(haloalkyl)-2-hydroxyacetonitrile such as 2-(3-chloropropyl)-2-hydroxy-2-methylacetonitrile; 2-(1-(protected amino)alkyl)-2-hydroxyacetonitrile such as 2-(1-alkoxycarbonylamino)-2-cyclohexylethyl)-2-hydroxyacetonitrile; 2-alkylthioalkyl-2-hydroxyacetonitrile such as 2-hydroxy-2-(2-methylthioethyl)acetonitrile; and 2-acyl-2-hydroxyacetonitrile such as 2-hydroxy-2-pivaloilacetonitrile.

A divalent group represented by R^1 and R^2 is not particularly limited, and the examples include an alkylene group containing 2 to 22 carbon atoms, norbornane-2-ylidene and 2-norbornene-5-ylidene.

The cyanohydrin used for the first aspect of the present invention can be obtained by known methods, i.e. by allowing alkali cyanide to act on a corresponding carbonyl compound or its sodium hydrogensulfite adduct. In a case of the use of cyanohydrin of the above formula (I), wherein R^1 and R^2 are different from each other, any of a (S)-form and an (R) form can be used, and this compound can be produced by, for example, such methods as an optical resolution of cyanohydrin obtained by acting alkali cyanide with a corresponding carbonyl compound or its sodium hydrogensulfite adduct; a method of asymmetricly adding hydrogen cyanide to a corresponding carbonyl compound in the presence of enzymes such as (S)-hydroxynitrilelyase and (R)-hydroxynitrilelyase extracted from plants (e.g. *Synthesis*, July 1990, 575-578; *Tetrahedron Letters*, 32, 2605-2608 (1991); Japanese Patent Application Filed Nos. 63-219388, 5-317065 and 9-227488); a method of asymmetricly hydrolyzing a material obtained by chemical synthesis of racemic cyanohydrinester, using enzymes (e.g.

Japanese Patent Application Filed No. 62-65688); and a method of asymmetricly adding hydrogen cyanide to a corresponding carbonyl compound in the presence of enzymes produced by a gene recombinant microorganism, into which a gene of enzymes such as (S)-hydroxynitrilelyase and (R)-hydroxynitrilelyase is incorporated (e.g. WO98/30711, Japanese Patent Application Filed No. 9-227488).

Mineral acid is preferably used for the hydrolysis in the first aspect of the present invention. The examples of such mineral acid include hydrochloric acid, sulfuric acid, nitric acid, boracic acid, phosphoric acid and perchloric acid, hydrochloric acid being preferable.

If an overfull usage of hydrolytic catalyst is used to cyanohydrin, it would be economically disadvantageous and lead to the decrease of the yield. However, the extremely low usage of catalyst would lead to a slow and insufficient reaction, or, in the use of optically active cyanohydrin as a material, it would lead to the reduction in the optical purity of the α -hydroxycarboxylic acid of interest. Accordingly, the preferable amount of hydrolytic catalyst is generally 1.5 to 10 equivalents to cyanohydrin, preferably 1.5 to 8 equivalents, and more preferably 2 to 7 equivalents.

Hereinafter, the hydrocarbon solvent used in the first aspect of the present invention is described. The hydrocarbon solvent used in the first aspect of the present invention is not particularly limited, as long as it is an organic compound consisting of carbon atoms and hydrogen atoms only, and it exists as a liquid at the reaction temperature of hydrolysis and separates from an aqueous phase.

Such carbon solvents include any of a linear or branched chain hydrocarbon, a cyclic hydrocarbon with or without side chain, and a chain hydrocarbon wherein the above cyclic hydrocarbon group is substituted. In addition, these hydrocarbons may have an unsaturated bond in a molecule thereof. Some representatives of the above

hydrocarbon solvents are shown below.

The linear or branched chain hydrocarbon includes pentane, hexane, heptane, octane and these structural isomers, e.g. chain hydrocarbons containing 5 to 16 carbon atoms, such as 2-methylpentane and 3-methylpentane.

The cyclic hydrocarbon with or without side chain includes cyclopentane, cyclohexane and structural isomers thereof, e.g. saturated monocyclic hydrocarbons containing 6 to 16 carbon atoms, such as methylcyclopentane and methylcyclohexane, and aromatic hydrocarbons such as benzene, toluene, trimethylbenzene, o-xylene, m-xylene, p-xylene or an isomeric mixture of xylene.

Among these hydrocarbon solvents, benzene, toluene and p-xylene are preferable, and toluene is more preferable.

A mixed solvent consisting of the combination of at least two of the above hydrocarbon solvents may also be used.

In the production method of the first aspect of the present invention, the above cyanohydrin is hydrolyzed in the presence of the above hydrocarbon solvent. That is, the reaction is performed by adding a hydrocarbon solvent, water containing a hydrolytic catalyst and cyanohydrin as a material to a reaction vessel. When left at rest, the reaction solution divides into both a hydrocarbon solvent phase and an aqueous phase, so it is preferable to perform hydrolysis, while stirring the reaction solution appropriately.

The usage of a hydrocarbon solvent in the first aspect of the present invention is 10 to 200 weight % to cyanohydrin as a material, and preferably 20 to 100 weight %.

The amount of water in the reaction mixture at initiation of the reaction is preferably 7 to 50 equivalents to cyanohydrin, and more preferably 10 to 40 equivalents.

In the production method of the first aspect of the present invention, when the maximum temperature at hydrolysis exceeds 90°C, the generation of by-products and coloration increases, or the purity of α -hydroxycarboxylic acid of interest is reduced. On the other hand, a reaction temperature of 40°C or less leads to the reaction proceeding insufficiently, resulting in a decrease of the yield. Accordingly, the temperature of reaction solution at hydrolysis is preferably 40 to 90°C, and more preferably 50 to 80°C. In addition, the reaction time is preferably 1 to 24 hours, 2 to 12 hours being more preferable.


After completion of the reaction, the α -hydroxycarboxylic acid of interest is isolated from the reaction solution. If the reaction solution is left at rest after hydrolysis, it divides into both a hydrocarbon solvent phase containing colored substances and by-products, and an aqueous phase containing α -hydroxycarboxylic acid. So, at that moment, the hydrocarbon solvent phase is separated and removed from the reaction solution. Then, extraction is performed by adding an organic solvent such as ethyl acetate to the remaining aqueous phase, and if necessary, the organic phase is washed with water and the solvent removed from the organic phase, thereby obtaining high purity α -hydroxycarboxylic acid of interest.

In the production method of the first aspect of the present invention, impurities such as colored substances and by-products generated during hydrolysis are extracted from the aqueous phase to the hydrocarbon solvent phase, while the α -hydroxycarboxylic acid of interest remains in the aqueous phase, so that high purity α -hydroxycarboxylic acid can easily be obtained, while suppressing the coloration of the reaction solution (an aqueous phase) and α -hydroxycarboxylic acid.

09870483-060101

The method of the second aspect of the present invention is to prevent reduction of the yield and optical purity of optically active α -hydroxycarboxylic acid caused by the use of an alcoholic solvent, an ester solvent, an ethereal solvent and/or a carboxylic solvent used in the production process of optically active cyanohydrin. The process of producing optically active cyanohydrin is not particularly limited, as long as these solvents are used as reaction solvents.

An example of the production process of optically active cyanohydrin includes a production process of optically active cyanohydrin by asymmetrically adding hydrogen cyanide to a carbonyl compound shown in the following formula (II):



wherein

R^1 and R^2 are different from each other, independently representing a hydrogen atom or a monovalent hydrocarbon group containing at most 22 carbon atoms, and in the above hydrocarbon group, each of $-CH_2-$ and CH_2 in $-CH_3$ may be substituted with a carbonyl group, a sulfonyl group, $-O-$ or $-S-$, $=CH_2$ may be substituted with $=O$ or $=S$; or C-H in $-CH_2$, C-H in $-CH_3$, C-H in $>CH-$, C-H in $=CH-$ and C-H in $=CH_2$ may be substituted with N or C-halogen, or

R^1 and R^2 may together form an asymmetric divalent group,

in the above organic solvent(s), in the presence of enzymes such as (S)-hydroxynitrilelyase and (R)-hydroxynitrilelyase extracted from plants or enzymes produced by a gene recombinant microorganism, into which a gene of those enzymes is incorporated (e.g. *Synthesis*, July 1990, 575-578; *Tetrahedron Letters*, 32, 2605-2608 (1991); Japanese Patent Application filed Nos. 63-219388, 5-317065 and 9-227488; WO98/30711).

The monovalent hydrocarbon group containing at most 22 carbon atoms in the above formula (II) includes a linear or branched chain hydrocarbon group, a monocyclic hydrocarbon group with or without side chain, a polycyclic hydrocarbon group with or

without side chain, a spiro hydrocarbon group with or without side chain, a ring-assembled structural hydrocarbon group with or without side chain, or a chain hydrocarbon group with the above cyclic-hydrocarbon. It includes any saturated or unsaturated hydrocarbon group, with the exception of unsaturated hydrocarbon groups having an allene structure ($C=C=C$). The linear or branched chain hydrocarbon group includes, for example, saturated chain hydrocarbon groups such as a linear alkyl group containing at least 1 carbon atom and a branched alkyl group containing at least 3 carbon atoms; unsaturated chain hydrocarbon groups such as a linear alkenyl group containing at least 2 carbon atoms, a branched alkenyl group containing at least 3 carbon atoms, a linear alkynyl group containing at least 3 carbon atoms, a branched alkynyl group containing at least 4 carbon atoms, a linear alkadienyl group containing at least 4 carbon atoms and a branched alkadienyl group containing at least 5 carbon atoms. The monocyclic hydrocarbon group includes, for example, saturated monocyclic hydrocarbon groups such as a cycloalkyl group without side chain which contains at least 3 carbon atoms and a cycloalkyl group with side chain which contains at least 4 carbon atoms in total; unsaturated monocyclic hydrocarbon groups such as a cycloalkenyl group without side chain which contains at least 4 carbon atoms, a cycloalkynyl group with side chain which contains at least 5 carbon atoms in total, a cycloalkadienyl group without side chain which contains at least 5 carbon atoms and a cycloalkadienyl group with side chain which contains at least 6 carbon atoms in total. The unsaturated monocyclic or polycyclic hydrocarbon group includes an aromatic hydrocarbon group including: an aromatic group without side chain which contains 6 to 22 carbon atoms in total such as a phenyl group, a 1-naphthyl group, a 2-naphthyl group and a 9-anthryl group; an aromatic group with side chain which contains at least 7 carbon atoms in total; a phenylphenyl group containing 12 carbon atoms and a phenylphenyl group with side chain which contains at least 13 carbon atoms in total, which are also included in a ring-assembled structural hydrocarbon group. The polycyclic hydrocarbon group includes a condensed cyclic hydrocarbon group without side chain which contains at least 6 carbon atoms, a condensed cyclic hydrocarbon

group with side chain which contains at least 7 carbon atoms in total, a bridged cyclic hydrocarbon group without side chain which contains at least 7 carbon atoms, a bridged cyclic hydrocarbon group with side chain which contains at least 8 carbon atoms in total, a spiro hydrocarbon group without side chain which contains at least 9 carbon atoms in total and a spiro hydrocarbon group with side chain which contains at least 10 carbon atoms in total. In addition, the above condensed cyclic hydrocarbon group without side chain includes those which contain at least 9 carbon atoms in total when one of its condensed rings is a benzene ring, and the above condensed cyclic hydrocarbon group with side chain includes those which contain at least 10 carbon atoms in total when one of its condensed rings is benzene ring. The ring-assembled structural hydrocarbon group includes a cycloalkyl-cycloalkyl group without side chain which contains at least 6 carbon atoms in total, a cycloalkyl-cycloalkyl group with side chain which contains at least 7 carbon atoms in total, a cycloalkylidene-cycloalkyl group without side chain which contains at least 6 carbon atoms in total and a cycloalkylidene-cycloalkyl group with side chain which contains at least 7 carbon atoms in total. "A cyclic hydrocarbon with side chain" in these cyclic hydrocarbons, corresponds to one having a chain hydrocarbon group attached to its ring. The above chain hydrocarbon group having a cyclic hydrocarbon group includes a linear alkyl group which is substituted with an aromatic group without side chain and contains at least 7 carbon atoms in total, a linear alkyl group which is substituted with an aromatic group with side chain and contains at least 8 carbon atoms in total, a branched alkyl group which is substituted with an aromatic group without side chain and contains at least 9 carbon atoms in total, a branched alkyl group which is substituted with an aromatic group with side chain and contains at least 10 carbon atoms in total, a linear alkenyl group which is substituted with an aromatic group without side chain and contains at least 8 carbon atoms in total, a linear alkenyl group which is substituted with an aromatic group with side chain and contains at least 9 carbon atoms in total, a branched alkenyl group which is substituted with an aromatic group without side chain and contains at least 9 carbon atoms in total, a branched alkenyl group which is substituted with an aromatic group with side chain

and contains at least 10 carbon atoms in total, a linear alkynyl group which is substituted with an aromatic group without side chain and contains at least 8 carbon atoms in total, a linear alkynyl group which is substituted with an aromatic group with side chain and contains at least 9 carbon atoms in total, a branched alkynyl group which is substituted with an aromatic group without side chain and contains at least 10 carbon atoms in total, a branched alkynyl group which is substituted with an aromatic group with side chain and contains at least 11 carbon atoms in total, a linear alkadienyl group which is substituted with an aromatic group without side chain and contains at least 10 carbon atoms in total, a linear alkadienyl group which is substituted with an aromatic group with side chain and contains at least 11 carbon atoms in total, a branched alkadienyl group which is substituted with an aromatic group without side chain and contains at least 11 carbon atoms in total, a branched alkadienyl group which is substituted with an aromatic group with side chain and contains at least 12 carbon atoms in total, a linear alkyl group which is substituted with a cycloalkyl group without side chain and contains at least 4 carbon atoms in total, a linear alkyl group which is substituted with a cycloalkyl group with side chain and contains at least 5 carbon atoms in total, a branched alkyl group which is substituted with a cycloalkyl group without side chain and contains at least 6 carbon atoms in total, a branched alkyl group which is substituted with a cycloalkyl group with side chain and contains at least 7 carbon atoms in total, a linear alkenyl group which is substituted with a cycloalkyl group without side chain and contains at least 5 carbon atoms in total, a linear alkenyl group which is substituted with a cycloalkyl group with side chain and contains at least 6 carbon atoms in total, a branched alkenyl group which is substituted with a cycloalkyl group without side chain and contains at least 6 carbon atoms in total, a branched alkenyl group which is substituted with a cycloalkyl group with side chain and contains at least 7 carbon atoms in total, a linear alkynyl group which is substituted with a cycloalkyl group without side chain and contains at least 5 carbon atoms in total, a linear alkynyl group which is substituted with a cycloalkyl group with side chain and contains at least 6 carbon atoms in total, a branched alkynyl group which is substituted with a cycloalkyl

group without side chain and contains at least 7 carbon atoms in total, a branched alkynyl group which is substituted with a cycloalkyl group with side chain and contains at least 8 carbon atoms in total, a branched alkadienyl group which is substituted with a cycloalkyl group without side chain and contains at least 8 carbon atoms in total, a branched alkadienyl group which is substituted with a cycloalkyl group with side chain and contains at least 9 carbon atoms in total.

Hereinafter, each of an aromatic group without side chain, an aromatic group with side chain, and a phenylphenyl group or a phenylphenyl group with side chain, is referred as an aryl group, and a linear or branched alkyl group substituted with the aryl group or groups is referred as an aralkyl group. Other cyclic hydrocarbon groups referring to both one having side chains on its ring and one having no side chains, are simply referred as, for example, cycloalkyl groups, unless otherwise specified. Also, chain hydrocarbon groups referring to both linear one and branched one are simply referred as, for example, alkyl groups.

When $-\text{CH}_2-$ in the above hydrocarbon group is substituted with a carbonyl group, sulfonyl group, $-\text{O}-$ or $-\text{S}-$, a ketone, sulfone, ether or thioether structure is introduced therein, respectively. When $-\text{CH}_2-$ in $-\text{CH}_3$ is substituted with a carbonyl group, $-\text{O}-$ or $-\text{S}-$, it converts into a formyl (aldehyde) group, a hydroxyl group or a mercapto group, respectively. When a terminal $=\text{CH}_2$ is substituted with $=\text{O}$ or $=\text{S}$, a ketone or thioketone structure is introduced therein. When $\text{C}-\text{H}$ in $-\text{CH}_2-$ is substituted with N , it converts into $-\text{NH}-$. When $\text{C}-\text{H}$ in $>\text{CH}-$ is substituted with N , it converts into $>\text{N}-$. When $\text{C}-\text{H}$ in $=\text{CH}-$ is substituted with N , it converts into $=\text{N}-$. When $\text{C}-\text{H}$ in a terminal $-\text{CH}_3$ is substituted with N , $-\text{NH}_2$ is introduced therein. When $\text{C}-\text{H}$ in $=\text{CH}_2$ is substituted with N , it converts into $=\text{NH}$. Further, each $\text{C}-\text{H}$ in $-\text{CH}_3$, $-\text{CH}_2-$, $=\text{CH}-$, $\equiv\text{CH}$ or $>\text{CH}-$ is substituted with a C-halogen, the carbon is substituted with a halogen atom. The substitution of carbon chains with $-\text{O}-$, $-\text{S}-$ or N corresponds to oxa-, thia- or aza-substitution of the hydrocarbon group, respectively. For example,

when these substitution take place in a ring carbon of the hydrocarbon ring, the hydrocarbon ring converts into a heterocyclic ring respectively containing oxygen, sulfur or nitrogen. The substitution of CH_2 and C-H in the hydrocarbon group may independently take place and it may further take place when CH_2 or C-H still remains on the carbon after the prior substitution. Further, the above substitutions may bring the conversion of $-\text{CH}_2-\text{CH}_3$ into $-\text{CO}-\text{O}-\text{H}$, a carboxylic acid structure.

A halogen atom, herein, refers to a fluorine atom, a chlorine atom, a bromine atom and an iodine atom, but a fluorine atom, a chlorine atom and a bromine atom are preferable.

Accordingly, the above hydrocarbon group may be selected from any chain hydrocarbon group and hydrocarbon group having ring-structured hydrocarbon group such as cyclic hydrocarbon groups, for example, saturated chain hydrocarbon groups such as linear or branched alkyl groups; unsaturated chain hydrocarbon groups such as linear or branched alkenyl groups, linear or branched alkynyl groups and linear or branched alkadienyl groups; saturated cyclic hydrocarbon groups such as cycloalkyl groups; unsaturated cyclic hydrocarbon groups such as cycloalkenyl groups, cycloalkynyl groups and cycloalkadienyl groups; and aromatic hydrocarbon groups such as aryl groups, aralkyl groups and arylalkenyl groups.

In more detail, the linear or branched alkyl groups include a methyl group, an ethyl group, a propyl group, an isopropyl group, a butyl group, a 1-methylpropyl group, a pentyl group, a 1-methylbutyl group, a hexyl group, a 1-methylpentyl group, a heptyl group, a 1-methylhexyl group, a 1-ethylpentyl group, an octyl group, a nonyl group, a decyl group, an undecyl group, a dodecyl group, a tridecyl group, a tetradecyl group, a 2-methylpropyl group, a 2-methylbutyl group, a 3-methylbutyl group, a 2-methylpentyl group, a 3-methylpentyl group, a 4-methylpentyl group, a methylhexyl group, a

methylheptyl group, a methyloctyl group, a methylnonyl group, a 1,1-dimethylethyl group, a 1,1-dimethylpropyl group, a 2,6-dimethylheptyl group, a 3,7-dimethyloctyl group and a 2-ethylhexyl group; cycloalkylalkyl groups include a cyclopentylmethyl group and a cyclohexylmethyl group; cycloalkyl groups include a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, a methylcyclopentyl group, a cyclohexyl group, a methylcyclohexyl group, a cycloheptyl group and a cyclooctyl group; and bicycloalkyl groups include a norbornyl group, a bicyclo [2.2.2] octyl group and an adamantyl group.

The linear or branched alkenyl groups include a vinyl group, an allyl group, a crotyl group (a 2-butenyl group) and an isopropenyl group (a 1-methylvinyl group); cycloalkenyl or cycloalkadienyl groups include a cyclopentenyl group, a cyclopentadienyl group, a cyclohexenyl group and a cyclohexadienyl group.

The linear or branched alkynyl groups include an ethynyl group, a propynyl group and a butynyl group. The aryl groups include a phenyl group, a 1-naphthyl group, a 2-naphthyl group, a 2-phenylphenyl group, a 3-phenylphenyl group, a 4-phenylphenyl group, a 9-anthryl group, a methylphenyl group, a dimethylphenyl group, a trimethylphenyl group, an ethylphenyl group, a methylethylphenyl group, a diethylphenyl group, a propylphenyl group and a butylphenyl group.

The aralkyl groups include a benzyl group, a 1-naphthylmethyl group, a 2-naphthylmethyl group, a phenethyl group (a 2-phenylethyl group), a 1-phenylethyl group, a phenylpropyl group, a phenylbutyl group, a phenylpentyl group, a phenylhexyl group, a methylbenzyl group, a methylphenethyl group, a dimethylbenzyl group, a dimethylphenethyl group, a trimethylbenzyl group, an ethylbenzyl group and a diethylbenzyl group.

The arylalkenyl groups include a styryl group, a methylstyryl group, an ethylstyryl group, a dimethylstyryl group and a 3-phenyl-2-propenyl group.

The above hydrocarbon groups, in which the CH_2 group is substituted with a carbonyl group, a sulfonyl group, O or S, or the C-H group is substituted with N or C-halogen, include groups having one or more structures such as ketone, aldehyde, carboxylic acid, sulfone, ether, thioether, amine, alcohol, thiol, halogen and heterocycles (e.g. an oxygen-containing heterocycle, a sulfur-containing heterocycle, a nitrogen-containing heterocycle, etc.) The oxygen-containing heterocycle, sulfur-containing heterocycle and nitrogen-containing heterocycle correspond to cyclic hydrocarbon groups in which their ring carbon is substituted with oxygen, sulfur and nitrogen, respectively. These heterocycles may contain two or more heteroatoms.

These substituted hydrocarbon groups may include a ketone structure such as an acetylmethyl group and an acetylphenyl group; a sulfone structure such as a methanesulfonylmethyl group; an ether structure such as a methoxymethyl group, a methoxyethyl group, an ethoxyethyl group, a methoxypropyl group, a butoxyethyl group, an ethoxyethoxyethyl group, a methoxyphenyl group, dimethoxyphenyl group and phenoxymethyl group; a thioether structure such as a methylthiomethyl group and a methylthiophenyl group; an amine structure such as an aminomethyl group, a 2-aminoethyl group, a 2-aminopropyl group, a 3-aminopropyl group, a 2,3-diaminopropyl group, a 2-aminobutyl group, a 3-aminobutyl group, a 4-aminobutyl group, a 2,3-diaminobutyl group, a 2,4-diaminobutyl group, a 3,4-diaminobutyl group, a 2,3,4-triaminobutyl group, a methylaminomethyl group, a dimethylaminomethyl group, a methylaminoethyl group, a propylaminomethyl group, a cyclopentylaminomethyl group, an aminophenyl group, a diaminophenyl group, an aminomethylphenyl group; an

oxygen-containing heterocycle such as a tetrahydrofuranlyl group, a tetrahydropyranlyl group and a morphorylethyl group; an oxygen-containing aromatic ring such as a furyl group, a furfuryl group, a benzofuryl group and a benzofurfuryl group; a sulfur-containing heterocycle such as a thienyl group; a nitrogen-containing aromatic ring such as a pyrrolyl group, an imidazolyl group, an oxazolyl group, a thiadiazolyl group, a pyridyl group, a pyrimidyl group, a pyridazinyl group, a pyrazinyl group, a tetrazinyl group, a quinolyl group, an isoquinolyl group, a pyridylmethyl group; an alcohol structure such as a 2-hydroxyethyl group, a 2-hydroxypropyl group, a 3-hydroxypropyl group, a 2,3-dihydroxypropyl group, a 2-hydroxybutyl group, a 3-hydroxybutyl group, a 4-hydroxybutyl group, a 2,3-dihydroxybutyl group, a 2,4-dihydroxybutyl group, a 3,4-dihydroxybutyl group, a 2,3,4-trihydroxybutyl group, a hydroxyphenyl group, a dihydroxyphenyl group, a hydroxymethylphenyl group and a hydroxyethylphenyl group; a thiol structure such as a 2-mercaptoethyl group, a 2-mercaptopropyl group, a 3-mercaptopropyl group, a 2,3-dimercaptopropyl group, a 2-mercaptobutyl group, a 3-mercaptobutyl group, a 4-mercaptobutyl group, a mercaptophenyl group; a halogenated hydrocarbon group such as a 2-chloroethyl group, a 2-chloropropyl group, a 3-chloropropyl group, a 2-chlorobutyl group, a 3-chlorobutyl group, a 4-chlorobutyl group, a fluorophenyl group, a chlorophenyl group, a bromophenyl group, a difluorophenyl group, a dichlorophenyl group, a dibromophenyl group, a chlorofluorophenyl group, a trifluorophenyl group, a trichlorophenyl group, a fluoromethylphenyl group and a trifluoromethylphenyl group; a compound having both an amine structure and an alcohol structure such as a 2-amino-3-hydroxypropyl group, a 3-amino-2-hydroxypropyl group, a 2-amino-3-hydroxybutyl group, a 3-amino-2-hydroxybutyl group, a 2-amino-4-hydroxybutyl group, a 4-amino-2-hydroxybutyl group, a 3-amino-4-hydroxybutyl group, a 4-amino-3-hydroxybutyl group, a 2,4-diamino-3-hydroxybutyl group, a 3-amino-2,4-dihydroxybutyl group, a 2,3-diamino-4-hydroxybutyl group, a 4-amino-2,3-dihydroxybutyl group, a 3,4-diamino-2-hydroxybutyl group, a 2-amino-3,4-

dihydroxybutyl group and an aminohydroxyphenyl group; a hydrocarbon group substituted with a halogen and a hydroxyl group such as a fluorohydroxyphenyl group, a chlorohydroxyphenyl group; and a carbon structure such as a carboxyphenyl group.

An asymmetric divalent group represented by R^1 and R^2 is not particularly limited, and the examples include norbornane-2-ylidene and 2-norbornene-5-ylidene.

The carbonyl compound shown in the above formula (II) includes: aromatic aldehyde such as benzaldehyde, m-phenoxybenzaldehyde, p-methylbenzaldehyde, o-chlorobenzaldehyde, m-chlorobenzaldehyde, p-chlorobenzaldehyde, m-nitrobenzaldehyde, 3,4-methylenedioxybenzaldehyde, 2,3-methylenedioxybenzaldehyde, phenylacetaldehyde and furfural; aliphatic aldehyde such as acetaldehyde, butylaldehyde, isobutylaldehyde, valeraldehyde and cyclohexanecarbaldehyde; saturated aliphatic ketone such as ethylmethylketone, butylmethylketone, methylpropylketone, isopropylmethylketone, methylpentylketone, methyl(2-methylpropyl)ketone and methyl(3-methylbutyl)ketone; unsaturated aliphatic ketone such as methyl(2-propenyl)ketone and (3-butenyl)methylketone; alkyl(haloalkyl)ketone such as (3-chloropropyl)methylketone; 2-(protected amino)aldehyde such as 2-(alkoxycarbonylamino)-3-cyclohexylpropionaldehyde; and alkylthio aliphatic aldehyde such as 3-methylthiopropionaldehyde.

In the production process of optically active cyanohydrin in the method of the second aspect of the present invention, a solvent comprising at least one organic solvent selected from a group consisting of an alcoholic solvent, an ester solvent, an ethereal solvent and a carboxylic solvent is used as a reaction solvent.

The alcoholic solvent includes aliphatic alcohol such as methanol, ethanol, propanol, isopropanol, 1-butanol, 2-butanol, isobutanol, t-butanol, hexanol, n-

amylalcohol and cyclohexanol as well as benzylalcohol and so on.


The ester solvent includes aliphatic ester such as methyl formate, methyl acetate, ethyl acetate, butyl acetate and methyl propionate.

The ethereal solvent includes aliphatic ether such as diethyl ether, dipropyl ether, diisopropyl ether, dibutyl ether, t-butylmethyl ether, dimethoxyethane and tetrahydrofuran.

The carboxylic solvent includes aliphatic carboxylic acid such as acetic acid.

Apart from the solvents listed above, the reaction solvent used in the second aspect of the present invention may also comprise a hydrocarbon-based solvent such as n-pentane, n-hexane, cyclohexane, benzene, toluene and xylene, and an aqueous buffer of pH7 or less such as a citrate buffer, a phosphate buffer and an acetate buffer.

In the production process of optically active cyanohydrin in the method of the second aspect of the present invention, the use of the carbonyl compound of the above formula (II) as a material results in the generation of optically active cyanohydrin corresponding to the above compound shown in the following formula (III):



wherein R¹ and R² have the same definitions as R¹ and R² in the above formula (II), and C* represents an asymmetric carbon atom. Subsequent to that, in the method of the

second aspect of the present invention, the as-is optically active cyanohydrin obtained in the first step, without isolation, is used for hydrolysis of the second step.

In the method of the second aspect of the present invention, an alcoholic solvent, an ester solvent, an ethereal solvent and/or a carboxylic solvent used as a reaction solvent in the first step were removed at that stage. Where a hydrocarbon-based solvent such as n-pentane, n-hexane, cyclohexane, benzene, toluene and xylene is used in combination with these organic solvent, the hydrocarbon-based solvent may not be removed.

Methods for removing the above solvent include, evaporation under an ordinary or reduced pressure, extraction with water and so on, but among these methods, evaporation under a reduced pressure is preferable in that this method is simple, having less influence on the second step.

Preferably, the removal of a reaction solvent in the first step of the second aspect of the present invention is preferably performed so that the amount of an alcoholic solvent, an ester solvent, an ethereal solvent and a carboxylic solvent in a reaction mixture subjected to hydrolysis is less than 10 weight %, and more preferably 5 weight %.

It is preferable to use mineral acid in performing hydrolysis in the second step of the second aspect of the present invention. Examples of mineral acid used herein include hydrochloric acid, sulfuric acid, nitric acid, boracic acid, phosphoric acid and perchloric acid, being hydrochloric acid preferable.

The preferable usage of mineral acid is 1 to 10 equivalents to optically active cyanohydrin which is contained in a reaction mixture subjected to hydrolysis. The usage of mineral acid of more than 10 equivalents to optically active cyanohydrin

would be economically disadvantageous, resulting in the decrease of yield. However, the usage of mineral acid of less than 1 equivalent to optically active cyanohydrin would lead to a slow and insufficient reaction, or the reduction in the optical purity of optically active α -hydroxycarboxylic acid of interest. The more preferable usage of mineral acid is 2 to 8 equivalents to optically active cyanohydrin.

Preferably, hydrolysis is performed under the condition of the maximum reaction temperature of 40 to 90°C. The maximum reaction temperature of more than 90°C would bring on the increase of by-products and coloration. On the other hand, the maximum reaction temperature of less than 40°C would result in the insufficient proceeding of reaction. Furthermore, in both cases, the optical purity of optically active α -hydroxycarboxylic acid of interest is reduced. The maximum reaction temperature is more preferably 40 to 80°C. Where the reaction temperature is less than 40°C, the reaction time is preferably set at 15 hours or less, more preferably 3 hours or less.

In hydrolysis reaction, a reaction solvent may be used, but such a solvent is not only particularly effective but may also reduce yield or optical purity, so it is preferable not to use solvents other than water. At the initiation of reaction, the amount of water contained in a reaction mixture including water contained in mineral acid used, is preferably 7 to 50 equivalents to optically active cyanohydrin, and more preferably 10 to 40 equivalents.

After reaction, to isolate α -hydroxycarboxylic acid of interest from the obtained reaction solution (which, at times, may become, a slurry), the solution is extracted with an organic solvent and is washed, as needed, and then the solvent is vaporized and exsiccated.


As stated above, by converting the cyano group of optically active

060101 060403 060703 060803 060903 061003 061103 061203 061303 061403 061503 061603 061703 061803 061903 062003 062103 062203 062303 062403 062503 062603 062703 062803 062903 063003 063103 063203 063303 063403 063503 063603 063703 063803 063903 064003 064103 064203 064303 064403 064503 064603 064703 064803 064903 065003 065103 065203 065303 065403 065503 065603 065703 065803 065903 066003 066103 066203 066303 066403 066503 066603 066703 066803 066903 067003 067103 067203 067303 067403 067503 067603 067703 067803 067903 068003 068103 068203 068303 068403 068503 068603 068703 068803 068903 069003 069103 069203 069303 069403 069503 069603 069703 069803 069903 070003 070103 070203 070303 070403 070503 070603 070703 070803 070903 071003 071103 071203 071303 071403 071503 071603 071703 071803 071903 072003 072103 072203 072303 072403 072503 072603 072703 072803 072903 073003 073103 073203 073303 073403 073503 073603 073703 073803 073903 074003 074103 074203 074303 074403 074503 074603 074703 074803 074903 075003 075103 075203 075303 075403 075503 075603 075703 075803 075903 076003 076103 076203 076303 076403 076503 076603 076703 076803 076903 077003 077103 077203 077303 077403 077503 077603 077703 077803 077903 078003 078103 078203 078303 078403 078503 078603 078703 078803 078903 079003 079103 079203 079303 079403 079503 079603 079703 079803 079903 080003 080103 080203 080303 080403 080503 080603 080703 080803 080903 081003 081103 081203 081303 081403 081503 081603 081703 081803 081903 082003 082103 082203 082303 082403 082503 082603 082703 082803 082903 083003 083103 083203 083303 083403 083503 083603 083703 083803 083903 084003 084103 084203 084303 084403 084503 084603 084703 084803 084903 085003 085103 085203 085303 085403 085503 085603 085703 085803 085903 086003 086103 086203 086303 086403 086503 086603 086703 086803 086903 087003 087103 087203 087303 087403 087503 087603 087703 087803 087903 088003 088103 088203 088303 088403 088503 088603 088703 088803 088903 089003 089103 089203 089303 089403 089503 089603 089703 089803 089903 090003 090103 090203 090303 090403 090503 090603 090703 090803 090903 091003 091103 091203 091303 091403 091503 091603 091703 091803 091903 092003 092103 092203 092303 092403 092503 092603 092703 092803 092903 093003 093103 093203 093303 093403 093503 093603 093703 093803 093903 094003 094103 094203 094303 094403 094503 094603 094703 094803 094903 095003 095103 095203 095303 095403 095503 095603 095703 095803 095903 096003 096103 096203 096303 096403 096503 096603 096703 096803 096903 097003 097103 097203 097303 097403 097503 097603 097703 097803 097903 098003 098103 098203 098303 098403 098503 098603 098703 098803 098903 099003 099103 099203 099303 099403 099503 099603 099703 099803 099903 100003 100103 100203 100303 100403 100503 100603 100703 100803 100903 101003 101103 101203 101303 101403 101503 101603 101703 101803 101903 102003 102103 102203 102303 102403 102503 102603 102703 102803 102903 103003 103103 103203 103303 103403 103503 103603 103703 103803 103903 104003 104103 104203 104303 104403 104503 104603 104703 104803 104903 105003 105103 105203 105303 105403 105503 105603 105703 105803 105903 106003 106103 106203 106303 106403 106503 106603 106703 106803 106903 107003 107103 107203 107303 107403 107503 107603 107703 107803 107903 108003 108103 108203 108303 108403 108503 108603 108703 108803 108903 109003 109103 109203 109303 109403 109503 109603 109703 109803 109903 110003 110103 110203 110303 110403 110503 110603 110703 110803 110903 111003 111103 111203 111303 111403 111503 111603 111703 111803 111903 112003 112103 112203 112303 112403 112503 112603 112703 112803 112903 113003 113103 113203 113303 113403 113503 113603 113703 113803 113903 114003 114103 114203 114303 114403 114503 114603 114703 114803 114903 115003 115103 115203 115303 115403 115503 115603 115703 115803 115903 116003 116103 116203 116303 116403 116503 116603 116703 116803 116903 117003 117103 117203 117303 117403 117503 117603 117703 117803 117903 118003 118103 118203 118303 118403 118503 118603 118703 118803 118903

The optically active cyanohydrin used as a material in the third aspect of the present invention is not particularly limited, as long as the cyanohydrin is an optically active substance (optically active α -hydroxynitrile) having, in a molecule thereof, at least one pair consisting of a hydroxyl group and a cyano group binding to an identical carbon atom. The optical purity of the above cyanohydrin is not particularly limited, as long as the purity is more than 80%, but an optical purity of 90 to 100% is preferable.

wherein

R^1 and R^2 are different from each other, independently representing a hydrogen atom, a



cycloalkynyl group with side chain which contains at least 5 carbon atoms in total, a cycloalkadienyl group without side chain which contains at least 5 carbon atoms and a cycloalkadienyl group with side chain which contains at least 6 carbon atoms in total. The unsaturated monocyclic or polycyclic hydrocarbon group includes an aromatic hydrocarbon group including: an aromatic group without side chain which contains 6 to 22 carbon atoms in total such as a phenyl group, a 1-naphthyl group, a 2-naphthyl group and a 9-anthryl group; an aromatic group with side chain which contains at least 7 carbon atoms in total; a phenylphenyl group containing 12 carbon atoms and a phenylphenyl group with side chain which contains at least 13 carbon atoms in total, which are also included in a ring-assembled structural hydrocarbon group. The polycyclic hydrocarbon group includes a condensed cyclic hydrocarbon group without side chain which contains at least 6 carbon atoms, a condensed cyclic hydrocarbon group with side chain which contains at least 7 carbon atoms in total, a bridged cyclic hydrocarbon group without side chain which contains at least 7 carbon atoms, a bridged cyclic hydrocarbon group with side chain which contains at least 8 carbon atoms in total, a spiro hydrocarbon group without side chain which contains at least 9 carbon atoms in total and a spiro hydrocarbon group with side chain which contains at least 10 carbon atoms in total. In addition, the above condensed cyclic hydrocarbon group without side chain includes those which contain at least 9 carbon atoms in total when one of its condensed rings is a benzene ring, and the above condensed cyclic hydrocarbon group with side chain includes those which contain at least 10 carbon atoms in total when one of its condensed rings is benzene ring. The ring-assembled structural hydrocarbon group includes a cycloalkyl-cycloalkyl group without side chain which contains at least 6 carbon atoms in total, a cycloalkyl-cycloalkyl group with side chain which contains at least 7 carbon atoms in total, a cycloalkylidene-cycloalkyl group without side chain which contains at least 6 carbon atoms in total and a cycloalkylidene-cycloalkyl group with side chain which contains at least 7 carbon atoms in total. "A cyclic hydrocarbon with side chain" in these cyclic hydrocarbons, corresponds to one having a chain hydrocarbon group attached to its ring. Such chain hydrocarbon group having a cyclic

hydrocarbon group includes a linear alkyl group which is substituted with an aromatic group without side chain and contains at least 7 carbon atoms in total, a linear alkyl group which is substituted with an aromatic group with side chain and contains at least 8 carbon atoms in total, a branched alkyl group which is substituted with an aromatic group without side chain and contains at least 9 carbon atoms in total, a branched alkyl group which is substituted with an aromatic group with side chain and contains at least 10 carbon atoms in total, a linear alkenyl group which is substituted with an aromatic group without side chain and contains at least 8 carbon atoms in total, a linear alkenyl group which is substituted with an aromatic group with side chain and contains at least 9 carbon atoms in total, a branched alkenyl group which is substituted with an aromatic group without side chain and contains at least 9 carbon atoms in total, a branched alkenyl group which is substituted with an aromatic group with side chain and contains at least 10 carbon atoms in total, a linear alkynyl group which is substituted with an aromatic group without side chain and contains at least 8 carbon atoms in total, a linear alkynyl group which is substituted with an aromatic group with side chain and contains at least 9 carbon atoms in total, a branched alkynyl group which is substituted with an aromatic group without side chain and contains at least 10 carbon atoms in total, a branched alkynyl group which is substituted with an aromatic group with side chain and contains at least 11 carbon atoms in total, a linear alkadienyl group which is substituted with an aromatic group without side chain and contains at least 10 carbon atoms in total, a linear alkadienyl group which is substituted with an aromatic group with side chain and contains at least 11 carbon atoms in total, a branched alkadienyl group which is substituted with an aromatic group without side chain and contains at least 11 carbon atoms in total, a branched alkadienyl group which is substituted with an aromatic group with side chain and contains at least 12 carbon atoms in total, a linear alkyl group which is substituted with a cycloalkyl group without side chain and contains at least 4 carbon atoms in total, a linear alkyl group which is substituted with a cycloalkyl group with side chain and contains at least 5 carbon atoms in total, a branched alkyl group which is substituted with a cycloalkyl group without side chain and contains at least 6 carbon

atoms in total, a branched alkyl group which is substituted with a cycloalkyl group with side chain and contains at least 7 carbon atoms in total, a linear alkenyl group which is substituted with a cycloalkyl group without side chain and contains at least 5 carbon atoms in total, a linear alkenyl group which is substituted with a cycloalkyl group with side chain and contains at least 6 carbon atoms in total, a branched alkenyl group which is substituted with a cycloalkyl group without side chain and contains at least 6 carbon atoms in total, a branched alkenyl group which is substituted with a cycloalkyl group with side chain and contains at least 7 carbon atoms in total, a linear alkynyl group which is substituted with a cycloalkyl group without side chain and contains at least 5 carbon atoms in total, a linear alkynyl group which is substituted with a cycloalkyl group with side chain and contains at least 6 carbon atoms in total, a branched alkynyl group which is substituted with a cycloalkyl group without side chain and contains at least 7 carbon atoms in total, a branched alkynyl group which is substituted with a cycloalkyl group with side chain and contains at least 8 carbon atoms in total, a branched alkadienyl group which is substituted with a cycloalkyl group without side chain and contains at least 8 carbon atoms in total, a branched alkadienyl group which is substituted with a cycloalkyl group with side chain and contains at least 9 carbon atoms in total.

Hereinafter, each of an aromatic group without side chain, an aromatic group with side chain, and a phenylphenyl group or a phenylphenyl group with side chain, is referred as an aryl group, and a linear or branched alkyl group substituted with the aryl group or groups is referred as an aralkyl group. Other cyclic hydrocarbon groups referring to both one having side chains on its ring and one having no side chains, are simply referred as, for example, cycloalkyl groups, unless otherwise specified. Also, chain hydrocarbon groups referring to both linear one and branched one are simply referred as, for example, alkyl groups.

When $-\text{CH}_2-$ in the above hydrocarbon group is substituted with a carbonyl

group, sulfonyl group, -O- or -S-, a ketone, sulfone, ether or thioether structure is introduced therein, respectively. When $-\text{CH}_2-$ in $-\text{CH}_3$ is substituted with a carbonyl group, -O- or -S-, it converts into a formyl (aldehyde) group, a hydroxyl group or a mercapto group, respectively. When a terminal $=\text{CH}_2$ is substituted with $=\text{O}$ or $=\text{S}$, a ketone or thioketone structure is introduced therein. When C-H in $-\text{CH}_2-$ is substituted with N, it converts into $-\text{NH}-$. When C-H in $>\text{CH}-$ is substituted with N, it converts into $>\text{N}-$. When C-H in $=\text{CH}-$ is substituted with N, it converts into $=\text{N}-$. When C-H in a terminal $-\text{CH}_3$ is substituted with N, $-\text{NH}_2$ is introduced therein. When C-H in $=\text{CH}_2$ is substituted with N, it converts into $=\text{NH}$. Further, each C-H in $-\text{CH}_3$, $-\text{CH}_2-$, $=\text{CH}-$, $\equiv\text{CH}$ or $>\text{CH}-$ is substituted with a C-halogen, the carbon is substituted with a halogen atom. The substitution of carbon chains with -O-, -S- or N corresponds to oxa-, thia- or aza-substitution of the hydrocarbon group, respectively. For example, when these substitution take place in a ring carbon of the hydrocarbon ring, the hydrocarbon ring converts into a heterocyclic ring respectively containing oxygen, sulfur or nitrogen. The substitution of CH_2 and C-H in the hydrocarbon group may independently take place and it may further take place when CH_2 or C-H still remains on the carbon after the prior substitution. Further, the above substitutions may bring the conversion of $-\text{CH}_2-\text{CH}_3$ into $-\text{CO}-\text{O}-\text{H}$, a carboxylic acid structure.

A halogen atom, herein, refers to a fluorine atom, a chlorine atom, a bromine atom and an iodine atom, but a fluorine atom, a chlorine atom and a bromine atom are preferable.

groups; unsaturated cyclic hydrocarbon groups such as cycloalkenyl groups, cycloalkynyl groups and cycloalkadienyl groups; and aromatic hydrocarbon groups such as aryl groups, aralkyl groups and arylalkenyl groups.

In more detail, the linear or branched alkyl groups include a methyl group, an ethyl group, a propyl group, an isopropyl group, a butyl group, a 1-methylpropyl group, a pentyl group, a 1-methylbutyl group, a hexyl group, a 1-methylpentyl group, a heptyl group, a 1-methylhexyl group, a 1-ethylpentyl group, an octyl group, a nonyl group, a decyl group, an undecyl group, a dodecyl group, a tridecyl group, a tetradecyl group, a 2-methylpropyl group, a 2-methylbutyl group, a 3-methylbutyl group, a 2-methylpentyl group, a 3-methylpentyl group, a 4-methylpentyl group, a methylhexyl group, a methylheptyl group, a methylloctyl group, a methylnonyl group, a 1,1-dimethylethyl group, a 1,1-dimethylpropyl group, a 2,6-dimethylheptyl group, a 3,7-dimethyloctyl group and a 2-ethylhexyl group; cycloalkylalkyl groups include a cyclopentylmethyl group and a cyclohexylmethyl group; cycloalkyl groups include a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, a methylcyclopentyl group, a cyclohexyl group, a methylcyclohexyl group, a cycloheptyl group and a cyclooctyl group; and bicycloalkyl groups include a norbornyl group, a bicyclo [2.2.2] octyl group and an adamantyl group.

The linear or branched alkenyl groups include a vinyl group, an allyl group, a crotyl group (a 2-butenyl group) and an isopropenyl group (a 1-methylvinyl group); cycloalkenyl or cycloalkadienyl groups include a cyclopentenyl group, a cyclopentadienyl group, a cyclohexenyl group and a cyclohexadienyl group.

The linear or branched alkynyl groups include an ethynyl group, a propynyl group and a butynyl group. The aryl groups include a phenyl group, a 1-naphthyl

group, a 2-naphthyl group, a 2-phenylphenyl group, a 3-phenylphenyl group, a 4-phenylphenyl group, a 9-anthryl group, a methylphenyl group, a dimethylphenyl group, a trimethylphenyl group, an ethylphenyl group, a methylethylphenyl group, a diethylphenyl group, a propylphenyl group and a butylphenyl group.

The aralkyl groups include a benzyl group, a 1-naphthylmethyl group, a 2-naphthylmethyl group, a phenethyl group (a 2-phenylethyl group), a 1-phenylethyl group, a phenylpropyl group, a phenylbutyl group, a phenylpentyl group, a phenylhexyl group, a methylbenzyl group, a methylphenethyl group, a dimethylbenzyl group, a dimethylphenethyl group, a trimethylbenzyl group, an ethylbenzyl group and a diethylbenzyl group.

The arylalkenyl groups include a styryl group, a methylstyryl group, an ethylstyryl group, a dimethylstyryl group and a 3-phenyl-2-propenyl group.

The above hydrocarbon groups, in which the CH_2 group is substituted with a carbonyl group, a sulfonyl group, O or S, or the C-H group is substituted with N or C-halogen, include groups having one or more structures such as ketone, aldehyde, carboxylic acid, sulfone, ether, thioether, amine, alcohol, thiol, halogen and heterocycles (e.g. an oxygen-containing heterocycle, a sulfur-containing heterocycle, a nitrogen-containing heterocycle etc.) The oxygen-containing heterocycle, sulfur-containing heterocycle and nitrogen-containing heterocycle correspond to cyclic hydrocarbon groups in which their ring carbon is substituted with oxygen, sulfur and nitrogen, respectively. These heterocycles may contain two or more heteroatoms.

These substituted hydrocarbon groups may include a ketone structure such as an acetylmethyl group and an acetylphenyl group; a sulfone structure such as a

methanesulfonylmethyl group; an ether structure such as a methoxymethyl group, a methoxyethyl group, an ethoxyethyl group, a methoxypropyl group, a butoxyethyl group, an ethoxyethoxyethyl group, a methoxyphenyl group, dimethoxyphenyl group and phenoxyethyl group; a thioether structure such as a methylthiomethyl group and a methylthiophenyl group; an amine structure such as an aminomethyl group, a 2-aminoethyl group, a 2-aminopropyl group, a 3-aminopropyl group, a 2,3-diaminopropyl group, a 2-aminobutyl group, a 3-aminobutyl group, a 4-aminobutyl group, a 2,3-diaminobutyl group, a 2,4-diaminobutyl group, a 3,4-diaminobutyl group, a 2,3,4-triaminobutyl group, a methylaminomethyl group, a dimethylaminomethyl group, a methylaminoethyl group, a propylaminomethyl group, a cyclopentylaminomethyl group, an aminophenyl group, a diaminophenyl group, an aminomethylphenyl group; an oxygen-containing heterocycle such as a tetrahydrofuran group, a tetrahydropyran group and a morphoylethyl group; an oxygen-containing aromatic ring such as a furyl group, a furfuryl group, a benzofuryl group and a benzofurfuryl group; a sulfur-containing heterocycle such as a thienyl group; a nitrogen-containing aromatic ring such as a pyrrolyl group, an imidazolyl group, an oxazolyl group, a thiazolyl group, a pyridyl group, a pyrimidyl group, a pyridazinyl group, a pyrazinyl group, a tetrazinyl group, a quinolyl group, an isoquinolyl group, a pyridylmethyl group; an alcohol structure such as a 2-hydroxyethyl group, a 2-hydroxypropyl group, a 3-hydroxypropyl group, a 2,3-dihydroxypropyl group, a 2-hydroxybutyl group, a 3-hydroxybutyl group, a 4-hydroxybutyl group, a 2,3-dihydroxybutyl group, a 2,4-dihydroxybutyl group, a 3,4-dihydroxybutyl group, a 2,3,4-trihydroxybutyl group, a hydroxyphenyl group, a dihydroxyphenyl group, a hydroxymethylphenyl group and a hydroxyethylphenyl group; a thiol structure such as a 2-mercaptoethyl group, a 2-mercaptopropyl group, a 3-mercaptopropyl group, a 2,3-dimercaptopropyl group, a 2-mercaptobutyl group, a 3-mercaptobutyl group, a 4-mercaptobutyl group, a mercaptophenyl group; a halogenated hydrocarbon group such as a 2-chloroethyl group, a 2-chloropropyl group, a 3-

chloropropyl group, a 2-chlorobutyl group, a 3-chlorobutyl group, a 4-chlorobutyl group, a fluorophenyl group, a chlorophenyl group, a bromophenyl group, a difluorophenyl group, a dichlorophenyl group, a dibromophenyl group, a chlorofluorophenyl group, a trifluorophenyl group, a trichlorophenyl group, a fluoromethylphenyl group and a trifluoromethylphenyl group; a compound having both an amine structure and an alcohol structure such as a 2-amino-3-hydroxypropyl group, a 3-amino-2-hydroxypropyl group, a 2-amino-3-hydroxybutyl group, a 3-amino-2-hydroxybutyl group, a 2-amino-4-hydroxybutyl group, a 4-amino-2-hydroxybutyl group, a 3-amino-4-hydroxybutyl group, a 4-amino-3-hydroxybutyl group, a 2,4-diamino-3-hydroxybutyl group, a 3-amino-2,4-dihydroxybutyl group, a 2,3-diamino-4-hydroxybutyl group, a 4-amino-2,3-dihydroxybutyl group, a 3,4-diamino-2-hydroxybutyl group, a 2-amino-3,4-dihydroxybutyl group and an aminohydroxyphenyl group; a hydrocarbon group substituted with a halogen and a hydroxyl group such as a fluorohydroxyphenyl group, a chlorohydroxyphenyl group; and a carbon structure such as a carboxyphenyl group.

The cyanohydrin shown in the above formula (IV) includes: 2-aryl-2-hydroxyacetonitrile such as mandelonitrile (2-hydroxy-2-phenylacetonitrile), 3-phenoxymandelonitrile (2-hydroxy-2-(3-phenoxyphenyl)acetonitrile), 4-methylmandelonitrile (2-hydroxy-2-(p-tolyl)acetonitrile), 2-chloromandelonitrile (2-(2-chlorophenyl)-2-hydroxyacetonitrile), 3-chloromandelonitrile (2-(3-chlorophenyl)-2-hydroxyacetonitrile), 4-chloromandelonitrile (2-(4-chlorophenyl)-2-hydroxyacetonitrile), 3-nitromandelonitrile (2-hydroxy-2-(3-nitrophenyl)acetonitrile), 3,4-methylenedioxymandelonitrile (2-hydroxy-2-(3,4-methylenedioxyphenyl)acetonitrile), 2,3-methylenedioxymandelonitrile (2-hydroxy-2-(2,3-methylenedioxyphenyl)acetonitrile), 2-benzyl-2-hydroxyacetonitrile and 2-(2-furyl)-2-hydroxyacetonitrile; 2-alkyl-2-hydroxyacetonitrile such as 2-hydroxy-2-methylacetonitrile, 2-hydroxy-2-propylacetonitrile, 2-hydroxy-2-isopropylacetonitrile, 2-butyl-2-hydroxyacetonitrile and 2-cyclohexyl-2-hydroxyacetonitrile; 2,2-dialkyl-2-

hydroxyacetonitrile such as 2-ethyl-2-hydroxy-2-methylacetonitrile, 2-butyl-2-hydroxy-2-methylacetonitrile, 2-hydroxy-2-methyl-2-propylacetonitrile, 2-hydroxy-2-isopropyl-2-methylacetonitrile, 2-hydroxy-2-methyl-2-pentylacetonitrile, 2-hydroxy-2-methyl-2-(2-methylpropyl)acetonitrile and 2-hydroxy-2-methyl-2-(3-methylbutyl)acetonitrile; 2-alkyl-2-alkenyl-2-hydroxyacetonitrile such as 2-hydroxy-2-methyl-2-(2-propenyl)acetonitrile and 2-(3-butenyl)-2-hydroxy-2-methylacetonitrile; 2-alkyl-2-(haloalkyl)-2-hydroxyacetonitrile such as 2-(3-chloropropyl)-2-hydroxy-2-methylacetonitrile; 2-(1-(protected amino)alkyl)-2-hydroxyacetonitrile such as 2-(1-alkoxycarbonylamino)-2-cyclohexylethyl)-2-hydroxyacetonitrile; 2-alkylthioalkyl-2-hydroxyacetonitrile such as 2-hydroxy-2-(2-methylthioethyl)acetonitrile; and 2-acyl-2-hydroxyacetonitrile such as 2-hydroxy-2-pivaloilacetonitrile.

An asymmetric divalent group represented by R^1 and R^2 is not particularly limited, as long as the above groups can convert a carbon atom, to which the groups bind, into an asymmetric carbon, and the examples include norbornane-2-ylidene and 2-norbornene-5-ylidene.

The optically active cyanohydrin used in the third aspect of the present invention may be any of a (S)-form and an (R) form, and this compound can be produced by, for example, such methods as an optical resolution of cyanohydrin obtained by acting alkali cyanide with a corresponding carbonyl compound or its sodium hydrogensulfite adduct; a method of asymmetrically adding hydrogen cyanide to a corresponding carbonyl compound in the presence of enzymes such as (S)-hydroxynitrilelyase and (R)-hydroxynitrilelyase extracted from plants (e.g. *Synthesis*, July 1990, 575-578; *Tetrahedron Letters*, 32, 2605-2608 (1991); Japanese Patent Application filed Nos. 63-219388, 5-317065 and 9-227488); a method of asymmetrically hydrolyzing a material obtained by chemical synthesis of racemic cyanohydrinester, using enzymes (e.g. Japanese Patent Application filed No. 62-65688); and a method of asymmetrically adding hydrogen cyanide to a corresponding carbonyl compound in the

presence of enzymes produced by a gene recombinant microorganism, to which a gene of enzymes such as (S)-hydroxynitrilelyase and (R)-hydroxynitrilelyase is incorporated (e.g. WO98/30711, Japanese Patent Application filed No. 9-227488).

The third aspect of the present invention is characterized in hydrolyzing optically active cyanohydrin, using at most 10 equivalents of mineral acid relative to the optically active cyanohydrin under the condition that maximum temperature when reacting is less than 90°C.

Examples of mineral acid used in the third aspect of the present invention include hydrochloric acid, sulfuric acid, nitric acid, boric acid, phosphoric acid and perchloric acid, hydrochloric acid being preferable.

The usage of more than 10 equivalents of mineral acid relative to optically active cyanohydrin would be economically disadvantageous, resulting in a decrease of yields. However, the extremely low usage of mineral acid would lead to a slow and insufficient reaction, or a reduction in the optical purity of the optically active α -hydroxycarboxylic acid of interest. The preferable usage of mineral acid is 1.5 to 8 equivalents relative to optically active cyanohydrin, and more preferably 2 to 7 equivalents.

A maximum reaction temperature of more than 90°C would bring on the increase of the generation of by-products and coloration, resulting in the reduction of the optical purity of optically active α -hydroxycarboxylic acid of interest. The maximum reaction temperature is preferably 40 to 80°C, more preferably 45 to 70°C. Where the reaction temperature is less than 40°C, the reaction time is preferably set at 15 hours or less, more preferably 3 hours or less.

A reaction solvent may be used, but such a solvent is not particularly

effective and may also reduce yield or optical purity, so it is preferable not to use solvents other than water. At initiation of the reaction, the amount of water contained in a reaction mixture including water in the mineral acid used, is preferably 7 to 50 equivalents relative to optically active cyanohydrin, and more preferably 10 to 40 equivalents.

After reaction, to isolate the α -hydroxycarboxylic acid of interest from the obtained reaction solution (which, at times, may be a slurry), the solution is extracted with an organic solvent and is washed, as needed, and then the solvent is vaporized and exsiccated.

The fourth aspect of the present invention is a method for producing optically active crystalline α -hydroxycarboxylic acid, which comprises crystallizing optically active α -hydroxycarboxylic acid in an aqueous solution. This crystallization may be carried out in the presence of a non-miscible organic solvent, and an example of such an organic solvent includes a hydrocarbon solvent such as benzene, toluene, and o-, m- and p-xylene, n-hexane, n-heptane, n-octane. Where the crystallization is carried out in the presence of a non-miscible organic solvent, the ratio between an aqueous solution and the organic solvent is preferably 1 : 0.05 — 1 : 1. An optically active crystalline α -hydroxycarboxylic acid is deposited by cooling the above optically active α -hydroxycarboxylic acid solution. The cooling temperature is not particularly limited, as long as the temperature is set at, or below, a temperature at which the optically active α -hydroxycarboxylic acid solution to be used becomes a saturated solution, but preferably at 30 °C or less, more preferably at 25 °C or less. It is better to perform the cooling as slowly as possible, preferably at a cooling rate of 0.5 °C/min or less.

As optically active α -hydroxycarboxylic acid to be used in the fourth aspect of the present invention, the optically active α -hydroxycarboxylic acid produced by the method of the first, second or third aspect of the present invention may be used. In

This specification includes part or all of the contents as disclosed in the specifications of Japanese Patent Applications Nos. 2000-166382, 2000-171186 and 2000-350695, which are the bases of the priority claim of the present application.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention is further described in the following examples. The example is provided for illustrative purposes only, and is not intended to limit the scope of the invention.

(Example 1.)

25.2g (0.150mol) of (R)-2-chloromandelonitrile whose optical purity is 98% e.e., 46.9g of 35% hydrochloric acid (HCl 0.450mol) and 7.56g of toluene were placed into a 100ml flask and stirred at room temperature for 1 hour, followed by reaction with heating at 70°C for 4 hours. After cooling to room temperature and leaving at rest, the upper phase, toluene was removed. 50 g of ethyl acetate was added to the remaining aqueous phase and mixed in a separatory funnel, while shaking, followed by separation of the organic phase from the water phase. The water phase was extracted with 50g of ethyl acetate, and the obtained organic phase was mixed with the previous organic phase, then the mixture was washed with 15g of water. The thus obtained solution containing the (R)-2-chloromandelic acid was exsiccated under a reduced pressure, and then washed with 25g of toluene and dried to obtain 26.9g of (R)-2-chloromandelic acid. The optical purity was 96.6% (e.e.) The yield of the obtained (R)-2-chloromandelic acid was 94.4%.

(Example 2.)

The same reaction as in Example 1 was carried out to produce (R)-2-chloromandelic acid, with the exception that the amount of toluene was set at 25.2g.

(Example 3.)

The same reaction as in Example 1 was carried out to produce (R)-2-chloromandelic acid, with the exception that benzene was used instead of toluene.

(Example 4.)

The same reaction as in Example 1 was carried out to produce (R)-2-chloromandelic acid, with the exception that p-xylene was used instead of toluene.

(Control example 1.)

The same reaction as in Example 1 was carried out to produce (R)-2-chloromandelic acid, with the exception that toluene was not added.

The results of Examples 1 to 4 and Control example 1 are shown in the following Table 1.

Table 1.

No.	Hydrocarbon solvents (ratio of weight to cyanohydrin)	Coloration of reaction solution	Yield of R-2CMA (%)	Purity of carboxylic acid (%)	Optical purity (% e.e.)
Example1	toluene (0.3)	light orange	94.4	99.0	96.6
Example2	toluene (1.0)	extremely light yellow	93.6	99.3	96.7
Example3	benzene (0.3)	light orange	94.1	99.1	96.6
Example4	p-xylene (0.3)	light orange	93.8	98.8	96.4
Control 1	none	dark reddish brown	93.6	96.5	96.2

R-2CMA: (R)-2-chloromandelic acid

(Preparation example 1.) Preparation of S-hydroxynitrilelyase

(Preparation of S-hydroxynitrilelyase)

According to genetic engineering procedures, S-hydroxynitrilelyase was prepared using a yeast *saccharomyces cerevisiae* as a host. That is to say, according to standard techniques, total mRNA was extracted from the leaf of *Manihot utilissima*. Then, using the obtained mRNA as a template, cDNA synthesis was carried out to produce cDNA. Meanwhile, based on the sequence of S-hydroxynitrilelyase gene from *Manihot utilissima* described in a reference (*Arch. Biochem. Biophys.* 311, 496-502 (1994)), the following primers were synthesized. With regard to the synthesized DNA of each of the following SEQ ID NOS: 1 and 2, a sequence listing is provided.

Sense primer: ggggaattcatggtactgcacattttgtctgattc (SEQ ID NO: 1)

Antisense primer: ggggtcgacctcaggattagaagccgccg (SEQ ID NO: 2)

With the synthesized primers, PCR (90°C, 30 seconds; 55°C, 30 seconds; 72°C, 60 seconds; 35 cycles in total) was carried out using the above cDNA as a template, to obtain a S-hydroxynitrilelyase gene. The analysis of the gene sequence showed that the sequence of the S-hydroxynitrilelyase gene matched with the sequence shown in the reference.

Subsequently, the obtained PCR fragments were inserted between the promoter and terminator of an expression vector, YEp352-GAP, to obtain a yeast episomic expression vector, YEp352-GC. This vector was transformed to a yeast *saccharomyces cerevisiae* Inv-Sc1 strain according to standard techniques, and then the obtained strains were cultured on a minimum selective medium, which did not contain uracil, so that recombinant yeast YEp352-GC-S2 strains containing an expression vector YEp352-GC were obtained.

Then, the obtained recombinant yeast YEp352-GC-S2 strains were cultured in a YNBDCas liquid medium (6.7g/L Yeast nitrogen base without amino acid (Difco),

20g/L glucose, 20g/L casamino acid, 40mg/mL L-tryptophan) for 24 hours to produce S-hydroxynitrilelyase in a cell thereof. Cells were collected from a recombinant cell culture solution by centrifugation, and then the obtained cell bodies were crushed by bead mill. The crushed cell bodies-containing solution was centrifuged to prepare a crude enzyme solution. The solution was roughly purified by ammonium sulfate fractionation, and the obtained solution was used as a S-hydroxynitrilelyase solution for the following experiments.

(Immobilization of S-hydroxynitrilelyase)

The above-prepared S-hydroxynitrilelyase was immobilized to Micro Bead Silica Gel 300A (Fuji Silysia Chemical Ltd.) The immobilization of enzyme was carried out by adding 200g of carrier to 1.0L of S-hydroxynitrilelyase solution (activity: 64U/ml, 0.02M HEPES-Na buffer (pH6.0)), stirring the mixture at 4°C for 24 hours, and performing the adsorption immobilization of enzyme protein to the carrier.

(1st step: Synthesis of optically active cyanohydrin with immobilized enzyme)

(Example 1')

200g of immobilized enzyme, 1.2L of t-butylmethyl ether (tBME) saturated with 10mM phosphate buffer (pH5.5), 127.2g (2.0mol) of benzaldehyde and 49.2g (3.0mol) of hydrocyanic acid were placed into a 2L-volume flask, and stirred at 20°C for 1 hour. After reaction, the amount and optical purity of the obtained cyanohydrin were determined by analyzing the reaction solution by HPLC.

(Example 2')

The same reaction as in Example 1' was carried out, with the exception that ethyl acetate was used as a solvent, instead of tBME.

(Example 3')

5.0 L of 25 weight % methanol solution containing 0.1M benzaldehyde, 0.3M

hydrocyanic acid and 0.05M sodium acetate, and 50g of immobilized enzyme were placed into a 5L-volum flask, followed by the same reaction as in Example 1'.

(Reference example)

The same reaction as in Example 1' was carried out, with the exception that hexane was used as a solvent, instead of tBME.

The results of the above reactions are shown in Table 2.

Table 2.

	Solvent	Invert ratio (%)	Optical purity (% ee)
Example 1'	tBME	98	more than 99.9
Example 2'	ethyl acetate	97	more than 99.9
Example 3'	methanol/water	96	99.8
Reference example	hexane	96	99.8

(2nd step: Hydrolysis)

(Example 1A)

After the reaction solution of Example 1' was filtrated to remove immobilized enzyme, 2.3g of para-toluene sulfonate monohydrate was added thereto and shaken. After that, one quarter of the obtained reaction solution (which corresponds to 0.5mol of initiation material) was concentrated with an evaporator until the amount of tBME became 2 weight %, and transferred into a 300mL flask with a reflux condenser. 156g of 35% hydrochloric acid (HCl, 1.5mol) was added thereto, and stirred at room temperature for 1 hour, followed by further stirring at 70°C for 5 hours. After cooling the mixture to room temperature, 175g of ethyl acetate was added thereto and mixed in a separatory funnel, while shaking, the organic phase was separated from water phase. Water phase was extracted with 175g of ethyl acetate again, and the obtained organic phase was mixed with the previous organic phase, then the mixture was washed with 50g of water. Such obtained organic phase was exsiccated under a reduced pressure,

(Control examples 1' to 3')

The same reaction as in Examples was carried out, with the exception that the amount of solvent remaining in the material for hydrolysis was changed.

The results of the hydrolysis reactions are shown in Table 3.

Table 3.

No.	Amount of solvent (wt %)	Yield of carboxylic acid (%)	Optical purity (% ee)
Example 1A	tBME (2)	95	99.0
Example 1B	tBME (5)	93	98.3
Example 2A	ethyl acetate (3)	94	98.8
Example 2B	ethyl acetate (8)	92	97.9
Example 3'	methanol (5)	95	97.1
Reference example	hexane (15)	96	98.5
Control example 1'	tBME (15)	86	89.0
Control example 2A	ethyl acetate (12)	89	86.4
Control example 2B	ethyl acetate (20)	85	84.0
Control example 3'	methanol (15)	90	92.5

Note: The above yield of carboxylic acid was determined by, first, obtaining the number of moles from the yield and HPLC purity, and then calculating the obtained value with the number of moles of cyanohydrin (including R-form) obtained in the 1st step.

(Example 1'')

25.2g (0.150mol) of (R) -2-chloromandelonitrile and 46.9g (HCl 0.450mol) of 35% hydrochloric acid were placed into a 100ml flask. After stirring at room temperature for 1 hour, the mixture was heated to 50°C followed by reaction for 15 hours. After cooling the reactant to room temperature, 50g of ethyl acetate was added thereto and mixed in a separatory funnel, while shaking, thereby separating the organic phase from aqueous phase. The aqueous phase was extracted with 50g of ethyl acetate, and the obtained organic phase was mixed with the previous organic phase, then the mixture was washed with 15g of water. The thus obtained solution containing (R) -2-

chloromandelic acid of interest was exsiccated under a reduced pressure, and then washed with 25g of toluene and dried to obtain 26.5g of (R)-2-chloromandelic acid. The purity of (R) -2-chloromandelic acid by HPLC was 99.0%, and the optical purity was 97.0% (e.e.) The yield of the obtained (R)-2-chloromandelic acid was 94%, and the packing density was 0.49.

(Examples 2'' and 3'', and Control examples 1'' and 2'')

(R)-2-chloromandelic acid was produced by the same method as described in Example 1'', with the exception that the amount of 35% hydrochloric acid used, reaction temperature and reaction time were changed.

The reaction conditions and results are shown in Table 4.

Each of the maximum reaction temperature in Examples 1'' to 3'' and Control examples 1'' and 2'' was 50°C, 80°C, 50°C, 100°C and 50°C, respectively.

Table 4.

No.	Acid (Usage molar ratio)	Reaction temperature & Reaction time	Reaction yield (%) of carboxylic acid	Purity (%) of carboxylic acid	Optical purity (% e.e.)	(R)-2- CIMA* yield (%)
Example 1''	35% HCl (3.0)	RT* 1h 50°C 15h	98	99.0	97.0	94
Example 2''	35% HCl (3.0)	RT* 1h 80°C 3h	97	98.5	92.6	87
Example 3''	35% HCl (6.0)	RT* 1h 50°C 12h	98	99.3	98.0	94
Control example 1''	35% HCl (3.0)	RT* 1h 100°C 2h	93	93.2	59.5	70
Control example 2''	35% HCl (20)	RT* 1h 50°C 12h	96	99.0	96.6	68

* RT: room temperature

*(R)-2-CIMA: (R)-2-chloromandelic acid

(Crystallization example)

Hydrolysis was carried out at 5 times the scale of Example 1''. That is, 126g (0.75mol) of (R)-2-chloromandelonitrile and 234.5g (HCl 2.25mol) of 35% hydrochloric acid were placed into a 50ml flask, and the mixture was stirred at room temperature for 1 hour. Then, the mixture was heated to 50°C, and reaction was performed for 15 hours.

After completion of the reaction, 120g of water was added to the reaction solution, followed by stirring at 50°C for 10 minutes to homogenize the solution. This solution was divided into 5 equal portions at a temperature of 50°C, then crystallization was carried out for each portion under different conditions. Each crystallization example is shown below.

(Example 1S)

The reaction solution, into which water was added, was cooled to 20°C at

constant cooling rate for 90 minutes followed by keeping at 20°C for 2 hours, and then the solution was filtrated. The obtained crystal was washed with 10g of water and then with 15g of toluene, followed by drying, thereby obtaining 25.3g of (R)-2-chloromandelic acid. The purity of (R)-2-chloromandelic acid was determined to be 99.3% by HPLC, the optical putiry being 98.7% (ee). The yield of the obtained (R)-2-chloromandelic acid was 89.0%. The packing density of the obtained (R)-2-chloromandelic acid was determined to be 0.58 by the method described later.

(Example 2S)

The same reaction as in Example 1S was carried out, with the exception that cooling was initiated after adding 15g of toluene. The yield of (R)-2-chloromandelic acid was 24.6g. The purity of (R)-2-chloromandelic acid was 99.6%, the optical purity being 99.5% (ee). The yield of the obtained (R)-2-chloromandelic acid was 87.2%, the packing density being 0.62.

(Example 3S)

The same reaction as in Example 2S was carried out, with the exception that cooling time was set at 30 minutes. The yield of (R)-2-chloromandelic acid was 24.8g. The purity of (R)-2-chloromandelic acid was 99.4%, the optical purity being 99.3% (ee). The yield of the obtained (R)-2-chloromandelic acid was 87.7%, the packing density being 0.60.

(Example 4S)

The same reaction as in Example 2S was carried out, with the exception that the reaction solution was cooled to 5°C. The yield of (R)-2-chloromandelic acid was 24.9g. The purity of (R)-2-chloromandelic acid was 99.3%, the optical purity being 99.1% (ee). The yield of the obtained (R)-2-chloromandelic acid was 88.0%, the packing density being 0.63.

(Example 5S)

The same reaction as in Example 2S was carried out, with the exception that *p*-xylene was used instead of toluene. The yield of (R)-2-chloromandelic acid was 24.8g. The purity of (R)-2-chloromandelic acid was 99.4%, the optical purity being 99.1% (ee). The yield of the obtained (R)-2-chloromandelic acid was 87.6%, the packing density being 0.62.

(Determination of packing density)

About 20g of crystal was naturally dropped through a 20mm-diameter funnel into a 50mL graduated cylinder, and the packing density of the crystal was calculated by measuring the weight and volume of the crystal.

Table 5.

	hydrocarbon	Final temperature (°C)	cooling time (min)	Yield of products (%)	Purity (%)	Optical purity (% ee)	Filling density (g/cm ³)
Example 1S	none	20	90	89.0	99.3	98.7	0.58
Example 2S	toluene	20	90	87.2	99.6	99.5	0.62
Example 3S	toluene	20	30	87.7	99.4	99.3	0.60
Example 4S	toluene	5	90	88.0	99.3	99.1	0.63
Example 5S	<i>p</i> -xylene	20	90	87.6	99.4	99.1	0.62

All the publications, patents and patent applications cited herein are incorporated herein by reference in their entirety.